



BEE'S FOAMING GEL

PHASE	INGREDIENTS	PROVIDER	INCI	(wt-%)
A	WATER		Water	45,90
	DIAPOK K-SF	NOF Corporation	Aqua, Sodium Methyl Cocoyl Taurate	26,70
	NISSAN ANON BDF-R	NOF Corporation	Aqua, Cocamido Propyl Betaine	16,70
	MACBIOBRIDE MG-120TIS	NOF Corporation	PEG-120 Methyl Glucose Triisostearate	2,50
	AMITOSE R-PD	Seiwa Kasei	Aqua, Dihydroxypropyl Arginine HCl, Pentyleneglycol	1,00
B	DN-P conc. 25	Dainihon Kasei	Cocamido Propyl Betaine, Cocamide MEA, Glycol Distearate, Sodium Chloride, Phenoxyethanol, Citric Acid, Water	4,00
	CITRIC ACID		Citric Acid	0,10
	FERMENTED HONEY PG	Katakura &Co-op Agri	Aqua, Propanediol, Gluconobacter/Honey Ferment Filtrate	2,00
	ROYAL BIOCYTE PX	Katakura &Co-op Agri	Aqua, Hydrolyzed Royal Jelly Protein, Phenoxyethanol	0,50
	PHENOXYETHANOL		Phenoxyethanol	0,50
	TEA LILY IV	Kobayashi Perfumery	Fragrance	0,10
			Total	100,00

- 1/ Add Phase A into a main vessel, and stir at 80°C.
- 2/ Cool down to room temperature.
- 3/ Add Phase B to Phase A and stir at room temperature.
- 4/ Fill the bulk in a suitable package.